

---

**Specialized tip/stalk-like and phalanx-like endothelial cells from embryonic stem cells.**

**Journal:** Stem Cells Dev

**Publication Year:** 2013

**Authors:** Alicia A Blancas, Lian E Wong, Drew E Glaser, Kara E McCloskey

**PubMed link:** 23249281

**Funding Grants:** Building Cardiac Tissue from Stem Cells and Natural Matrices

**Public Summary:**

Endothelial cells (EC) generated in vitro from stem cells are desirable for their potential in a variety of in vitro models and cell-based therapeutic approaches; however, EC can take on a number of functionally and phenotypically distinct specializations. Here, we show the generation of functionally distinct EC subpopulations, including (1) the pro-angiogenic migrating tip-like and proliferative stalk-like EC, and (2) the less migratory cobblestone-shaped phalanx-like EC. Both embryonic stem cell (ESC)-derived EC subpopulations are generated from outgrowths of Flk-1<sup>+</sup> vascular progenitor cells with high levels of vascular endothelial growth factor treatment, while the phalanx-like ESC-derived EC (ESC-EC) are subsequently isolated by selecting for cobblestone shape. Compared with the ESC-derived angiogenic endothelial cells (named ESC-AEC) that contain only 14% Flt-1<sup>+</sup> and 25% Tie-1<sup>+</sup> cells, the selected phalanx-like ESC-EC express higher numbers of cells expressing the phalanx markers Flt-1<sup>+</sup> and Tie-1<sup>+</sup>, 89% and 90%, respectively. The ESC-AEC also contain 35% CXCR4<sup>+</sup> tip cells, higher expression levels of stalk marker Notch-1, and lower expression levels of Tie-2 compared with the phalanx-type ESC-EC that do not contain discernible numbers of CXCR4<sup>+</sup> tip cells. Perhaps most notably, the ESC-AEC display increased cell migration, proliferation, and 3 times more vessel-like structures after 48 h on Matrigel compared with the phalanx-like ESC-EC. This work analyzes, for the first time, the presence of distinct EC subtypes (tip/stalk, and phalanx) generated in vitro from ESC, and shows that phalanx-like EC can be purified and maintained in culture separate from the tip/stalk-like containing EC.

**Scientific Abstract:**

Endothelial cells (EC) generated in vitro from stem cells are desirable for their potential in a variety of in vitro models and cell-based therapeutic approaches; however, EC can take on a number of functionally and phenotypically distinct specializations. Here, we show the generation of functionally distinct EC subpopulations, including (1) the pro-angiogenic migrating tip-like and proliferative stalk-like EC, and (2) the less migratory cobblestone-shaped phalanx-like EC. Both embryonic stem cell (ESC)-derived EC subpopulations are generated from outgrowths of Flk-1<sup>+</sup> vascular progenitor cells with high levels of vascular endothelial growth factor treatment, while the phalanx-like ESC-derived EC (ESC-EC) are subsequently isolated by selecting for cobblestone shape. Compared with the ESC-derived angiogenic endothelial cells (named ESC-AEC) that contain only 14% Flt-1<sup>+</sup> and 25% Tie-1<sup>+</sup> cells, the selected phalanx-like ESC-EC express higher numbers of cells expressing the phalanx markers Flt-1<sup>+</sup> and Tie-1<sup>+</sup>, 89% and 90%, respectively. The ESC-AEC also contain 35% CXCR4<sup>+</sup> tip cells, higher expression levels of stalk marker Notch-1, and lower expression levels of Tie-2 compared with the phalanx-type ESC-EC that do not contain discernible numbers of CXCR4<sup>+</sup> tip cells. Perhaps most notably, the ESC-AEC display increased cell migration, proliferation, and 3 times more vessel-like structures after 48 h on Matrigel compared with the phalanx-like ESC-EC. This work analyzes, for the first time, the presence of distinct EC subtypes (tip/stalk, and phalanx) generated in vitro from ESC, and shows that phalanx-like EC can be purified and maintained in culture separate from the tip/stalk-like containing EC.

---

**Source URL:** <https://www.cirm.ca.gov/about-cirm/publications/specialized-tipstalk-and-phalanx-endothelial-cells-embryonic-stem-cells>